

RESEARCH ARTICLE

Open Access

A randomised controlled phase II trial of pre-operative celecoxib treatment reveals anti-tumour transcriptional response in primary breast cancer

Rita D Brandão^{1,2†}, Jürgen Veeck^{2,3,4,5*†}, Koen K Van de Vijver^{2,4}, Patrick Lindsey¹, Bart de Vries^{2,4}, Catharina HMJ van Elssen^{6,7}, Marinus J Blok¹, Kristien Keymeulen^{2,8}, Torik Ayoubi⁹, Hubert JM Smeets¹, Vivianne C Tjan-Heijnen^{2,3} and Pierre S Hupперets^{2,3}

Abstract

Introduction: Cyclooxygenase-2 (COX-2) is frequently over-expressed in primary breast cancer. In transgenic breast cancer models, over-expression of COX-2 leads to tumour formation while COX-2 inhibition exerts anti-tumour effects in breast cancer cell lines. To further determine the effect of COX-2 inhibition in primary breast cancer, we aimed to identify transcriptional changes in breast cancer tissues of patients treated with the selective COX-2 inhibitor celecoxib.

Methods: In a single-centre double-blind phase II study, thirty-seven breast cancer patients were randomised to receive either pre-operative celecoxib (400 mg) twice daily for two to three weeks ($n = 22$) or a placebo according to the same schedule ($n = 15$). Gene expression in fresh-frozen pre-surgical biopsies (before treatment) and surgical excision specimens (after treatment) was profiled by using Affymetrix arrays. Differentially expressed genes and altered pathways were bioinformatically identified. Expression of selected genes was validated by quantitative PCR (qPCR). Immunohistochemical protein expression analyses of the proliferation marker Ki-67, the apoptosis marker cleaved caspase-3 and the neo-angiogenesis marker CD34 served to evaluate biological response.

Results: We identified 972 and 586 significantly up- and down-regulated genes, respectively, in celecoxib-treated specimens. Significant expression changes in six out of eight genes could be validated by qPCR. Pathway analyses revealed over-representation of deregulated genes in the networks of proliferation, cell cycle, extracellular matrix biology, and inflammatory immune response. The Ki-67 mean change relative to baseline was -29.1% ($P = 0.019$) and -8.2% ($P = 0.384$) in the treatment and control arm, respectively. Between treatment groups, the change in Ki-67 was statistically significant ($P = 0.029$). Cleaved caspase-3 and CD34 expression were not significantly different between the celecoxib-treated and placebo-treated groups.

Conclusions: Short-term COX-2 inhibition by celecoxib induces transcriptional programs supporting anti-tumour activity in primary breast cancer tissue. The impact on proliferation-associated genes is reflected by a reduction of Ki-67 positive cells. Therefore, COX-2 inhibition should be considered as a treatment strategy for further clinical testing in primary breast cancer.

Trial registration: ClinicalTrials.gov NCT01695226.

* Correspondence: juergen.veeck@rwth-aachen.de

† Contributed equally

²GROW - School for Oncology and Developmental Biology, Maastricht University Medical Centre[†], P. Debye laan 25, Maastricht, 6202 AZ, The Netherlands

Full list of author information is available at the end of the article

Introduction

Cancer development is associated with chronic immune activation, but the mechanisms behind this observation are not fully understood [1,2]. In addition, the inflammatory processes that follow tumour formation provide a microenvironment in which the development of malignant disease may be enhanced [3]. The involvement of chronic immune activation has been supported by several lines of evidence in which an association between non-steroidal anti-inflammatory drug (NSAID) consumption and decreased risk of cancer development, including breast cancer, has been demonstrated [4-11]. However, the exact mechanisms by which NSAIDs exert inhibiting effects on tumour development have not yet been completely elucidated.

Mediators of inflammatory responses, such as the cyclooxygenase (COX)-derived prostaglandins (PG), play an important role in tumour formation and provide a target for therapeutic intervention [3]. PGs have important functions in every organ system and regulate a variety of physiological functions, such as immunity, maintenance of vascular integrity and bone metabolism [12]. Elevated COX expression in breast cancer was first suggested by the finding of elevated PG production in breast cancer cells [13]. To date, two different COX genes have been characterised, *COX-1* and *COX-2*. *COX-1* is constitutively expressed by almost all human cells and, therefore, differs from *COX-2* expression which is normally absent but is inducible by a wide spectrum of growth factors, pro-inflammatory cytokines [14,15] and tumour-promoting compounds [16,17]. Consistently, *COX-2* is abundantly expressed in breast cancer tissue [18] and its enforced over-expression in mammary gland epithelia of transgenic mice results in breast tumour development [19], suggesting that *COX-2* might be an interesting therapeutic target in breast cancer.

While several pre-clinical studies have indeed shown anti-tumour capacities of *COX-2* inhibition, the treatment effects on primary breast cancer in the clinical setting remain elusive. Therefore, we aimed to determine for the first time transcriptional changes in primary breast cancer tissue of women with early breast cancer after treatment with the selective *COX-2* inhibitor celecoxib. In our randomised controlled trial we found that upon celecoxib treatment numerous genes are differentially expressed in breast cancer tissues with an overall anti-tumour activity, suggesting that *COX-2* inhibition should be further considered for clinical testing as a treatment option in breast cancer.

Methods

Patients and study design

The study was a double-blind, randomised, placebo-controlled phase II pre-surgical trial of celecoxib in early

breast cancer. Exclusion criteria were: HIV, hepatitis B virus (HBV) or hepatitis C virus (HCV) positivity, known hypersensitivity to NSAIDs, patients already using NSAIDs or systemic use of corticosteroids. Informed consent was obtained prior to entering the trial and the Medical Ethics Committee of the Maastricht University Medical Centre+ (MUMC+) approved the study. We estimated that to test 18,500 genes at the 5% significance level and ensure 80% power, 23 samples were needed to detect differentially expressed genes by t-tests with a fold change of at least 1.5 (<http://bioinformatics.mdanderson.org/MicroarraySampleSize/>). Initially, 45 patients were recruited between 2005 and 2007 and randomly allocated 2:1 to the treatment ($n = 30$) or placebo group ($n = 15$). Celecoxib was pre-surgically administered for two to three weeks at 400 mg twice daily, whereas patients in the control arm received a placebo on the same schedule. Eight patients allocated to the treatment arm dropped out because these patients were operated earlier, thus drug compliance was insufficient (Figure 1). Tumour histology was assessed according to criteria defined by the World Health Organization (WHO) [20], while staging was performed according to the Union for International Cancer Control (UICC) criteria [21]. Tumours were graded following the system of Bloom and Richardson, as modified by Elston and Ellis [22]. Patient characteristics are described in Table 1. Importantly, in our study design patients acted as their own control, with




Table 1 Clinico-pathological characteristics of breast cancer patients.

Variable	Categorisation	Control	Celecoxib
		Number = 15	Number = 22
Mean age ± SD (years) ^a		57 ± 12	51 ± 12
Histological tumour type ^b			
	Invasive ductal carcinoma	13 (87%)	19 (86%)
	Invasive lobular carcinoma	2 (13%)	3 (14%)
Tumour size ^c			
	pT1	10 (67%)	8 (36%)
	pT2	5 (33%)	13 (59%)
	pT3	0 (0%)	1 (5%)
Lymph node status ^c			
	Negative (pN0)	11 (73%)	10 (45%)
	Positive (pN1-3)	4 (27%)	12 (55%)
Tumour grade ^d			
	G1	5 (33%)	1 (5%)
	G2	4 (27%)	7 (32%)
	G3	6 (40%)	14 (63%)
Oestrogen receptor status ^e			
	ER positive	11 (73%)	19 (86%)
	ER negative	4 (27%)	3 (14%)
Progesterone receptor status ^e			
	PgR positive	10 (67%)	17 (77%)
	PgR negative	5 (33%)	5 (23%)
HER2 status ^f			
	HER2 positive	3 (20%)	3 (14%)
	HER2 negative	12 (80%)	18 (82%)
	Unknown	0 (0%)	1 (5%)

^aAt time of diagnosis; ^baccording to Tavassoli and Devilee [20]; ^caccording to Sobin and Wittekind [21]; ^daccording to Elston and Ellis [22]; ^epositivity defined as > 10% positive cells as analysed by immunohistochemistry [60]; ^fpositivity defined by immunohistochemistry (score 3+) or by dual-color FISH. Percentages may not sum-up to 100 due to rounding. FISH, fluorescence *in situ* hybridisation.

a direct comparison of the final surgical specimen with the initial diagnostic biopsy. The inclusion of a placebo group served to observe a possible confounding impact of the disease and the experimental procedure, thereby allowing determination of a differential impact of celecoxib only.

Biopsy method

Before patient allocation to the groups, two to three core needle biopsies from the centre of the primary tumour were obtained using a 14-gauge needle under ultrasound guidance. At surgery, a central sample of the excised tumour was obtained. One part of the biopsies and surgical excision specimens was snap-frozen in liquid nitrogen within 30 minutes after removal and stored at -80°C until use, while the remaining part was formalin-fixed and paraffin-embedded (FFPE). All specimens subjected to gene expression analyses had a tumour cellularity of at least 80%, as determined by haematoxylin and eosin stained sections.

RNA isolation, cRNA production and fragmentation, array hybridisation and scanning

After homogenisation of fresh-frozen tissue specimens, total RNA was isolated using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol. DNase treatment was performed with the RNase-Free DNase Set (Qiagen, Hilden, Germany) followed by purification of the RNA samples using the RNeasy Mini-kit (Qiagen). RNA quantity and purity were determined spectrophotometrically using the Nanodrop ND-1000 (Nanodrop Technologies, Wilmington, DE, USA) and RNA integrity was assessed by determining the RNA 28S/18S ratio using the Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA). Biotinylated aRNA was synthesised and fragmented using the GeneChip IVT Express Kit from Affymetrix (Santa Clara, CA, USA). Hybridisation to Affymetrix Human Genome U133 Plus 2.0 arrays and subsequent scanning was performed following the manufacturer's guidelines using the GeneChip scanner 3000 (Affymetrix). Microarray datasets are publicly available at ArrayExpress database [23] under accession number E-MTAB-566.

Microarray data analysis

Images of the arrays were quantified with GCOS software (Affymetrix). The chip description file (CDF) used for the analysis was an update created and freely distributed by the microarray lab of the University of Michigan [24] based on Ensembl (version 10). The treatment effects on global gene expression were assessed by microarray analyses of subjects after celecoxib treatment versus subjects before treatment and controls (before and after treatment). Thus, expression changes have been adjusted for differences in the pre- and post-treatment specimens of the control group. The genes were analysed using a Gaussian linear regression including the hybridisation and labelling spikes and COX-2. The inference criterion used for comparing the models is their ability to predict the observed data, that is, models are compared directly through their minimised minus log-likelihood. When the numbers of parameters in models differ, they are penalised by adding the number of estimated parameters, a form of the Akaike information criterion (AIC) [25]. For each gene, the group effect only after intervention was then added to the model. The gene under consideration was found to be differentially expressed if the AIC decreased compared to the model not containing this effect. All statistical analyses presented were performed using the freely available program R [26] and the publicly available library 'growth' [27].

Validation of microarrays by quantitative RT-PCR

Quantitative RT-PCR (qPCR) was performed for selected genes on all 74 samples in order to validate the

results obtained in the microarray study. As a reference, TATA box binding protein (*TBP*) and RNA, 18S ribosomal 1 (*RN18S1*) were included as housekeeping genes. The list of genes and the primers that were used are described in Additional file 1, Table S1. Excess biotinylated aRNA (before the fragmentation step) was used for validation, after cDNA synthesis with M-MuLV Reverse Transcriptase (Finnzymes, Espoo, Finland) and oligo (dT), (Invitrogen) using the SensiMix SYBR Kit (Quantace, London, UK) following the manufacturer's protocol. qPCRs were run on the 7900HT system (Applied Biosystems, Singapore). Results were analysed using a Gaussian linear regression similar to microarray data. Expression of housekeeping genes (*TBP* and *GAPDH*) and *COX-2* were included during the analysis. The AIC was used to assess whether there was a difference between the controls and patients (group effect).

Analysis of functional categories

Significantly altered genes that were found to have a fold-change difference of at least 10% were classified into categories of biological processes and molecular functions using DAVID (Database for Annotation, Visualisation and Integrated Discovery) [28] and PathVisio [29]. DAVID and PathVisio analyses for pathway enrichment were performed first for all significantly altered genes, then for up- and down-regulated genes separately. DAVID results are listed together with *P*-values corrected with the Bonferroni method, as this was the most conservative method used by the software. A *P*-value of 0.05 was used as a cut-off value. In both analyses we filtered the pathways or gene ontology (GO) terms for which less than five and more than 150 genes were found, as those were either too specific or too general for our analysis. In PathVisio, the gene database Hs_Derby_20110601.bridge and the pathway collection from WikiPathways [30] were used to obtain a ranked list of pathways with differentially expressed genes. PathVisio results were sorted by Z-score, which is the standard statistical test under the hypergeometric distribution. Only pathways with a Z-score above 3, which corresponds to *P*-values of 0.0013 or lower, were selected.

Tissue marker expression

Biomarkers of response were assessed by immunohistochemistry on FFPE tissues. Cell proliferation was assessed using the MIB1 mouse monoclonal antibody to Ki-67 [31]. Measurement of apoptosis was performed by cleaved caspase-3 (Asp175) staining [32]. Tumour cells/section were counted for Ki-67 (1,000 cells) and apoptotic index (3,000 cells). The apoptotic index was expressed as a percentage of the number of cells displaying apoptotic bodies, scoring 0 if < 0.5%, +1 if between 0.5% and 2%, and +3 if > 2%. Samples were also assessed for expression of CD34 [33] using the Chalkley method [34].

Statistical analyses

Ki-67 values at baseline and time of surgery were expressed as geometric mean proportions of the baseline and transformed into percentage changes. Ki-67 changes within groups were compared using a paired t-test, differences in Ki-67 and CD34 between groups by using an unpaired t-test, and the Mann-Whitney test was used for group comparisons. Differences in caspase-3 expression were assessed by the Freeman-Halton extension of the Fisher's exact probability test. All tests were performed two-sided at the 5% significance level.

Results

Celecoxib treatment changes gene expression in breast carcinoma tissue

Two to three weeks of celecoxib treatment significantly altered the expression of 1,558 genes in breast cancer tissues, of which 972 genes were up- and 586 genes were down-regulated after treatment and adjustment to control tissue gene expression. The 50 most strongly up-regulated and down-regulated genes are presented in Additional file 2, Table S2 and Additional file 3, Table S3, respectively. For all of the selected genes, except two, significant expression changes were confirmed by qPCR (Figure 2).

The list of the biological pathways, in which significantly changed genes were over-represented, identified by DAVID analysis, is shown in Table 2. Biological processes identified by PathVisio are shown in Table 3. In both approaches down-regulated genes were consistently over-represented in cell cycle-related processes and proliferation. Up-regulated genes were over-represented in extracellular matrix (ECM) organisation, cell adhesion, and blood vessel development in DAVID analyses, whereas PathVisio results suggested an implication of further tumour suppressive pathways, for example, complement




Figure 2 qPCR validation of selected genes differentially expressed in celecoxib-treated samples as determined by microarray analysis. Fold-change and the 95% CI (error bars) are shown. Expression of six out of eight genes analysed (indicated by asterisks) was significantly changed in agreement with the microarray analysis.

Table 2 Over-represented functional categories within genes up- and/or down-regulated after celecoxib treatment using DAVID.

GO-ID	Term	Count	Fold enrichment	P-value ^a
Up-regulated				
0030198	Extracellular matrix organisation	28	4.657	6.35E-08
0009611	Response to wounding	70	2.337	1.32E-07
0007155	Cell adhesion	83	2.108	3.24E-07
0022610	Biological adhesion	83	2.105	3.46E-07
0001944	Vasculature development	43	2.951	1.20E-06
0001568	Blood vessel development	42	2.953	1.96E-06
0043062	Extracellular structure organisation	33	3.502	2.35E-06
0051270	Regulation of cell motion	34	3.047	4.89E-05
0001501	Skeletal system development	46	2.510	5.23E-05
0030334	Regulation of cell migration	31	3.173	9.15E-05
0040012	Regulation of locomotion	33	2.973	1.51E-04
0042127	Regulation of cell proliferation	83	1.834	2.27E-04
0042060	Wound healing	32	2.944	3.14E-04
0007160	Cell-matrix adhesion	20	3.931	1.42E-03
0048514	Blood vessel morphogenesis	33	2.692	1.56E-03
0031589	Cell-substrate adhesion	21	3.745	1.57E-03
0006928	Cell motion	53	1.984	7.97E-03
0001525	Angiogenesis	25	2.922	1.15E-02
0009612	Response to mechanical stimulus	14	4.324	3.91E-02
Down-regulated				
0006259	DNA metabolic process	52	3.018	5.10E-09
0007049	Cell cycle	64	2.411	1.85E-07
0006260	DNA replication	28	4.387	3.98E-07
0022402	Cell cycle process	50	2.569	4.12E-06
0000278	Mitotic cell cycle	37	2.898	3.70E-05
0000279	M phase	32	2.832	6.92E-04
0051276	Chromosome organisation	40	2.461	7.18E-04
0006974	Response to DNA damage stimulus	34	2.656	1.20E-03
0000087	M phase of mitotic cell cycle	25	3.229	1.76E-03
0022613	Ribonucleoprotein complex biogenesis	22	3.544	2.07E-03
0000280	Nuclear division	24	3.156	4.53E-03
0007067	Mitosis	24	3.156	4.53E-03
0022403	Cell cycle phase	35	2.454	4.81E-03
0042254	Ribosome biogenesis	17	4.062	8.27E-03
0048285	Organelle fission	24	3.031	9.01E-03
0006281	DNA repair	27	2.785	9.53E-03
0006396	RNA processing	41	2.185	1.01E-02
0031396	Regulation of protein ubiquitination	15	4.434	1.30E-02
0031397	Negative regulation of protein ubiquitination	13	5.177	1.31E-02
0031145	Anaphase-promoting complex-dependent proteasomal ubiquitin-dependent protein catabolic process	12	5.461	2.01E-02
0016071	mRNA metabolic process	31	2.428	2.52E-02
0034470	ncRNA processing	20	3.133	4.28E-02
0051439	Regulation of ubiquitin-protein ligase activity during mitotic cell cycle	12	4.986	4.85E-02

^aP-values shown were corrected for multiple testing using the Bonferroni method; categories ranked by P-value; functional categories with P-values < 0.05 and at least five, but less than 150 genes meeting the criterion are shown.

Table 3 List of local pathways in PathVisio significantly affected by the celecoxib treatment.

MAPP Name	number changed	number measured	% changed	Z-score ^a
Up-regulated				
Complement and coagulation cascades	15	50	30.00%	5.65
Senescence and autophagy	18	100	18.00%	3.61
RANKL/RANK signalling pathway	11	50	22.00%	3.58
Complement activation, classical pathway	5	15	33.33%	3.55
Matrix metalloproteinases	7	29	24.14%	3.14
TGF beta signalling pathway	18	113	15.93%	3.03
Inflammatory response pathway	7	30	23.33%	3.03
Oxidative stress	6	24	25.00%	3.01
Adipogenesis	19	125	15.20%	2.89
Endochondral ossification	11	61	18.03%	2.82
TWEAK signalling pathway	8	40	20.00%	2.73
SREBP signalling	6	28	21.43%	2.56
Down-regulated				
Cell cycle	17	95	17.89%	6.62
DNA replication	9	41	21.95%	5.61
Nucleotide metabolism	5	17	29.41%	5.12
Proteasome degradation	10	57	17.54%	4.97
Electron transport chain	10	81	12.35%	3.61
G1 to S cell cycle control	7	67	10.45%	2.51

^aZ-score is the standard statistical test under the hypergeometric distribution; pathways ranked by Z-score. Z-score > 1.96 (corresponding to *P*-value = 0.05) and at least five, but less than 150 genes meeting the criterion are shown. TGF, tumour growth factor.

activation, senescence and autophagy, and transforming growth factor- β (TGF- β) signaling.

Effects of celecoxib on cell cycle gene expression

DAVID and PathVisio consistently identified biological processes referring to regulation of cell cycle and proliferation, in particular among those genes down-regulated after treatment. An important downstream DNA damage response gene, *GADD45A* (Growth arrest and DNA-damage-inducible, alpha), was significantly up-regulated (Figure 3). Consistent with a putative activation of the G2/M checkpoint and cell cycle arrest due to DNA damage response, target genes *CCNB1* (G2/mitotic-specific cyclin-B1) and *CCNB2* (G2/mitotic-specific cyclin-B2) were significantly down-regulated after treatment.


Effects of celecoxib on ECM degradation gene expression

The majority of the matrix metalloproteinase (MMP) family members have been associated with tumour progression. The conversion of pro-MMP to active MMP-2 requires membrane type MT1-MMP (MMP-14), a trans-membrane protein that is activated intracellularly by the convertase *FURIN* [35]. The down-regulation of the protein convertase *FURIN* in the celecoxib-treated group potentially leads to less activation of MT1-MMP. Additionally, the effect of MMP-2 on proteolysis was inhibited either by up-regulation of *TIMP1*, *TIMP2*, *TIMP3*, or by *RECK* (Figure 4). The up-regulated *RECK*

exerts inhibitory effects on the conversion of pro-MMP-2 to MMP-2 and on the activation of pro-MMP-9 to MMP-9. In summary, our data suggest that degradation of ECM proteins was significantly inhibited in the celecoxib-treated group.

Recruitment of tumour-infiltrating leukocytes to breast carcinoma tissue after celecoxib treatment

Breast cancer tissue of celecoxib-treated patients showed a significantly increased expression of MHC class II genes, including *HLA-DR α* and *HLA-DR β 2*, *CD74* (MHC class II invariant chain) and *HLA-DM*, but not *HLA-DQ* and *HLA-DOA* [see Additional file 4, Table S4]. MHC class I gene expression was not significantly changed. Subsequent to the increased expression of HLA-class II genes, co-stimulatory markers of antigen-presenting cells (*CD83*) and the monocyte differentiation antigen *CD14* were up-regulated after celecoxib treatment. Infiltration of antigen-presenting cells was supported by increased expression of the Pattern Recognition Receptors (PRR), Toll-like receptor-2 (*TLR2*) and MD-1 (*LY86*). Gene expression of the classical B cell markers, *CD20* and *CD19*, was not altered, although there was increased expression of immunoglobulin J chain. As myeloid-derived suppressor cells (MDSC) immune suppressive cells also express MHC class II molecules, and their presence correlates with COX-2 over-expression [36], induction of MDSC signalling was investigated. However, MDSC





induction seems unlikely, since the expression of MDSC-signalling genes *ARG1* and *NOS2* is not altered. Moreover, expression of important effector molecules, such as granzymes and perforin, was not affected. The increased infiltration of leukocytes observed in the breast tumour seems restricted to macrophages and dendritic cells.

Change of tissue biomarker expression

To confirm the transcriptional changes, we determined expression of protein markers for proliferation, apoptosis, and neo-angiogenesis. The proliferation marker Ki-67 was assessed on paired pre- and post-treatment tissues.

Due to a lack of further tissue, apoptotic marker cleaved caspase-3 and neo-angiogenesis marker CD34 were assessed only on post-treatment tissues. Baseline Ki-67 positivity in the control group (geometric mean: 10.0%; 95% CI: 5.5 to 18.3) was not significantly different from baseline Ki-67 positivity in the treatment group (geometric mean: 13.4%; 95% CI: 9.8 to 18.3) ($P = 0.915$). The change in Ki-67 is shown for individual patients according to treatment or control arm in Figure 5. The geometric mean change in Ki-67 relative to baseline in the treatment arm was -29.1% (95% CI: -40.1% to -16.2%, $P = 0.019$), whereas in the control arm it was -8.2% (95% CI:



-23.1 to 9.5%, $P = 0.384$). There was a significant change difference between the two groups ($P = 0.029$). In contrast, the apoptotic index was not significantly different in post-treatment tissues ($P = 0.231$). The number of CD34 positive cells was slightly higher in celecoxib-treated tissues, but this was statistically insignificant. The geometric means (95% CI) of the 'Chalkley mean' value were 6.8 (5.4 to 8.5) in the control group and 7.7 (6.8 to 8.8) in the treatment group ($P = 0.376$).

Discussion

In this study, we analysed the transcriptional changes seen in primary breast cancer tissue following short-term celecoxib treatment. To accomplish this, we used global gene expression profiles from paired pre- and post-treatment specimens. After adjustment to the control group, we identified a large number of differentially expressed genes after treatment that are involved in the regulation of cancer-associated pathways, such as cell cycle and proliferation, ECM biology, and inflammatory response, amongst others. Most convincingly, COX-2 inhibition induced gene expression patterns indicative of a decelerated cell cycle and reduced proliferation. Celecoxib may induce G2/M arrest by p53 activation,

leading to *GADD45A* up-regulation, which in turn inhibits cyclin-B1 and cyclin-B2 expression and promotes G2/M arrest (Figure 3). A G2/M arrest is mainly forced after DNA damage to enable the initiation of DNA repair mechanisms [37]. Our finding is in line with previous studies investigating the effects of celecoxib on cancerous cells *in vitro*. Dvory-Sobol and colleagues demonstrated that celecoxib induces G2/M arrest associated with cyclin-B1 down-regulation in *K-RAS*-transformed enterocytes [38], and in the COX-2 expressing murine breast cancer cell line MCa-35, celecoxib induced a G2/M arrest followed by apoptosis [39]. Interestingly, equally treated lung cancer A549 cells lacking COX-2 expression showed increased DNA damage, but low levels of apoptosis in these cells suggested a selective effect of celecoxib on COX-2 expressing cells [39]. Celecoxib seems to increase DNA damage in irradiated cells, enhancing their radiosensitivity [40]. However, the mechanisms behind increased DNA damage in celecoxib-treated tumour cells remains poorly understood. Our results are further consistent with a study of colorectal cancer (CRC) cell lines in which more than 1,000 genes were identified as differentially expressed after celecoxib treatment, clustering of which revealed significant changes of cell cycle control, apoptosis, and lymphocyte infiltration [41]. Also in a study on primary CRC, celecoxib-induced gene expression changes significantly interfered with proliferation pathways [42]. In summary, we have confirmed a positive treatment effect of COX-2 inhibition on cell proliferation-related transcriptional programs in primary breast carcinomas, as has been previously demonstrated by several *in vitro* and *in vivo* studies.

Disruption of the basement membrane is a hallmark of malignancy. Degradative enzymes, such as MMPs, are produced by tumour cells and by resident and infiltrating cells as a response to the tumour, and contribute to matrix degradation and facilitate tumour invasion. MMP-2, MMP-9, and other members of the MMP family have been associated with tumour progression [43]. In particular, MMP-2 and MMP-9 activity appears to be inhibited by celecoxib in our study, the first by up-regulation of MMP antagonists (*TIMP1*, *TIMP2*, and *TIMP3*), the latter by up-regulation of the MMP-9 inhibitor *RECK*. An involvement of selective COX-2 inhibition in matrix stability by decreasing MMP activity and tumour invasiveness has been previously demonstrated in breast and CRC cancer models [44,45], thus being in good agreement with our data.

Several lines of evidence demonstrated that immune cell infiltration in tumours is enhanced by celecoxib treatment, which is associated with a better prognosis [46,47]. In our study, increased infiltration of antigen-presenting cells is supported by gene expression data whereas other immune cells of both the innate and adaptive immune system do

not seem to be affected by celecoxib treatment. Most up-regulated genes within this category belong to MHC class II. Comparable data on MHC class I and II induction have been reported by Lönnroth *et al.* in CRC patients using a NSAID [47].

In order to investigate whether observed gene expression changes after COX-2 inhibition have translated to a biologically relevant effect, we analysed protein markers for proliferation, apoptosis, and neo-angiogenesis in primary tissues. Suppression of the proliferation marker Ki-67 has been previously reported as a surrogate marker for decreased aromatase activity in oestrogen receptor (ER) positive breast cancer treated with aromatase inhibitors (AI) for two weeks [48,49]. Since COX-2 expression is positively correlated with tumour aromatase content [50], we were interested whether COX-2 inhibition would also lead to a reduction in Ki-67 positivity. As expected from previous studies [51,52], Ki-67 was not significantly reduced in the control arm. In contrast, the celecoxib arm showed a significant suppression of Ki-67, confirming the reduced proliferation observed in our gene expression data and suggesting an indirect treatment effect on aromatase activity. Although the Ki-67 suppressive effect was only modest (-29%) as compared to the AI anastrozole (-75%) [49], it was similar to the Ki-67 suppression achieved with another AI, that is, raloxifene (-24%) [52]. Other than Ki-67, caspase-3 and CD34 were not significantly changed after celecoxib treatment, although this has to be interpreted cautiously due to the lack of baseline data. However, in a previous neo-adjuvant study in breast cancer, two weeks of celecoxib did not result in a biological response of proliferation and apoptosis, as determined by Ki-67 staining and TUNEL assays, respectively [53]. Notably, the referenced study analysed fewer patients and used half the drug dose that we applied. Taken together, we hypothesise that two weeks of COX-2 inhibition may not be sufficient to translate all transcriptional activation to a measurable biological phenotype. In particular, suppression of blood vessel development may take effect only after longer drug exposure, which should be taken into consideration when designing future clinical trials of COX-2 inhibition in cancer.

In breast cancer, COX-2 over-expression is positively associated with HER2 over-expression [54] and with tumour aromatase content [50]. Therefore, COX-2 inhibition might prove beneficial, especially in combination with trastuzumab in HER2 positive breast cancer or with AI in hormone receptor (HR) positive disease. In trastuzumab-refractory metastatic breast cancer COX-2 inhibition was previously shown to be inactive [55]. However, improved efficacy and endpoint benefits of celecoxib in combination with AI were reported in post-menopausal metastatic breast cancer, although these were pronounced only in tamoxifen resistant patients [56,57]. Although promising,

further studies are needed in order to elucidate a clinical benefit of COX-2 inhibition in combination with other drugs in breast cancer treatment.

The strength of our study is the trial design, which allows patients to act as their own control, and the inclusion of a placebo group, which served to exclude potentially confounding effects by the disease and the experimental procedures. It would be interesting to sub-analyse the gene expression and Ki-67 in stratified breast cancer subtypes, as for example, HR positive *versus* negative disease, but this is hindered by the low number of HR negative patients in the control ($n = 5$) and the treatment arm ($n = 1$). Of note, however, there is a trend of bias towards poorer prognostic factors in the celecoxib arm, which could not have been predicted or changed before the end of the treatment procedure. Although this bias must be considered a limitation of the study, we speculate that the observed anti-tumour transcriptional response in this arm may have been under-estimated, and thus could become even more evident by comparing clinically highly similar arms.

Unfortunately, eight patients dropped out of the study due to early surgery, all of them from the treatment arm, resulting in a reduced statistical power to detect differentially expressed genes. However, in our microarray analysis we found more differentially expressed genes than expected by chance alone (1,558 observed versus 920 expected). After identification of the differentially expressed genes, pathway analyses were performed in order to identify genes clustering within the same biological pathway, as those genes are very likely to be true positives.

Large meta-analyses have recently shown that the non-selective COX inhibitor aspirin is protective against cancer development. Regular intake of aspirin significantly reduced the risk of several cancers, including breast cancer (OR = 0.88) [58], and three years onwards of daily aspirin intake reduced cancer incidence in both women and men (OR = 0.75) [59]. These studies suggest that selective COX-2 inhibitors may have similar effects on cancer incidence albeit with the advantage of causing less adverse side effects associated with aspirin use, such as gastrointestinal bleeding. COX-1 is expressed constitutively in many different tissues, whereas COX-2 is conditionally induced, such as in inflammation, suggesting that selective COX-2 inhibition might prove more successful in cancer prevention than non-selective COX inhibitors.

Conclusions

Our clinical trial provides substantial evidence for an anti-tumour activity of celecoxib based on global transcriptional changes and suppression of Ki-67 protein in primary breast cancer tissues, encouraging further clinical trials of celecoxib or its derivative molecules in breast cancer. Our

study confirms results from previous *in vitro* and *in vivo* studies as we found a large number of cell-cycle and proliferation-associated genes to be differentially expressed in celecoxib-treated primary breast cancer tissues. Our aim was to determine the global transcriptional response to short-term COX-2 inhibition in primary tumours, and our results indicate that proliferation and ECM biology are significantly influenced, possibly underlying the proposed anti-tumour activity. Future trials of COX-2 inhibition considering other primary endpoints, such as pathological or clinical response, should take into account that effects of a transcriptional response may need a longer time to translate into a measurable clinical benefit.

Additional material

Additional file 1: Table S1 showing the primer sequences used for qPCR validation.

Additional file 2: Table S2 showing the 50 most significantly up-regulated genes after treatment.

Additional file 3: Table S3 showing the 50 most significantly down-regulated genes after treatment.

Additional file 4: Table S4 showing genes of the immune surveillance mechanism and their deregulation after treatment.

Abbreviations

AI: aromatase inhibitor; AIC: Akaike information criterion; COX-2: cyclooxygenase-2; CRC: colorectal cancer; DAVID: Database for Annotation, Visualisation and Integrated Discovery; ECM: extracellular matrix; ER: oestrogen receptor; FFPE: formalin-fixed and paraffin-embedded; GEP: gene expression profiling; GO: gene ontology; HR: hormone receptor; MDSC: myeloid-derived suppressor cells; MMP: matrix metalloproteinase; NSAID: non-steroidal anti-inflammatory drug; PG: prostaglandin; qPCR: quantitative polymerase chain reaction; RT-PCR: reverse transcriptase polymerase chain reaction.

Authors' contributions

RDB performed microarray data analyses, participated in qPCR validation, data interpretation and co-drafted the manuscript. JV participated in qPCR validation, data interpretation, statistical analyses of immunohistochemical results and co-drafted the manuscript. KKVdV performed pathological review and immunohistochemical scoring, participated in data interpretation and critically revised the manuscript. PL provided statistical analyses of microarray data and critically revised the manuscript. BdV provided support in immunohistochemical scoring and critically revised the manuscript. CHMJVE performed immunohistochemical staining, participated in data interpretation and critically revised the manuscript. MJB provided support for the data mining and critically revised the manuscript. KK provided surgical support in tissue acquisition and critically revised the manuscript. TA performed microarray analyses and critically revised the manuscript. HJMS participated in the conception and design of the study and critically revised the manuscript. VCTH participated in the conception and design of the study and critically revised the manuscript. PSH conceived and coordinated the study and critically revised the manuscript. All authors read and approved the final version of the manuscript.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

We are grateful to the patients who participated in this trial. The expert technical help of Sabina Vanherle is greatly appreciated. This work was

supported by a grant obtained by CORE Fund - 2002 (financed by Pharmacia & Upjohn; Pfizer since 2003) and the Fundação para a Ciência e Tecnologia (SFRH/BD/32386/2006 to R.D.B), co-financed by the POPH and European Community Fund.

Author details

¹Department of Clinical Genetics, Unit of Clinical Genomics, Maastricht University Medical Centre⁺, P. Debyealaan 25, Maastricht, 6229 HX, The Netherlands. ²GROW - School for Oncology and Developmental Biology, Maastricht University Medical Centre⁺, P. Debyealaan 25, Maastricht, 6202 AZ, The Netherlands. ³Department of Internal Medicine, Division of Medical Oncology, Maastricht University Medical Centre⁺, P. Debyealaan 25, Maastricht, 6229 HX, The Netherlands. ⁴Department of Pathology, Maastricht University Medical Centre⁺, P. Debyealaan 25, Maastricht, 6229 HX, The Netherlands. ⁵Institute of Pathology, RWTH Aachen University Hospital, Pauwelsstr. 30, Aachen, 52074, Germany. ⁶Department of Internal Medicine, Division of Haematology, Maastricht University Medical Centre⁺, P. Debyealaan 25, Maastricht, 6229 HX, The Netherlands. ⁷Pharmacell BV, Oxfordlaan 70, Maastricht, 6229 EV, The Netherlands. ⁸Department of Surgery, Maastricht University Medical Centre⁺, P. Debyealaan 25, Maastricht, 6202 AZ, The Netherlands. ⁹Saint James School of Medicine, Basic Science and Premedical Faculty, Kralendijk, Bonaire, Netherlands Antilles.

Received: 20 August 2012 Revised: 27 February 2013

Accepted: 3 April 2013 Published: 8 April 2013

References

- O'Byrne KJ, Dalgleish AG: **Chronic immune activation and inflammation as the cause of malignancy.** *Br J Cancer* 2001, **84**:473-483.
- Grivennikov SI, Greten FR, Karin M: **Immunity, inflammation, and cancer.** *Cell* 2010, **140**:883-899.
- Wang D, DuBois RN: **Eicosanoids and cancer.** *Nat Rev Cancer* 2010, **10**:181-193.
- Friedman G, Ury H: **Initial screening for carcinogenicity of commonly used drugs.** *J Natl Cancer Inst* 1980, **65**:723-733.
- Harris RE, Namboodiri KK, Farrar WB: **Nonsteroidal antiinflammatory drugs and breast cancer.** *Epidemiology* 1996, **7**:203-205.
- Thun MJ, Namboodiri MM, Heath CW: **Aspirin use and reduced risk of fatal colon cancer.** *N Engl J Med* 1991, **325**:1593-1596.
- Giardiello FM, Hamilton SR, Krush AJ, Piantadosi S, Hylind LM, Celano P, Booker SV, Robinson CR, Offerhaus GJ: **Treatment of colonic and rectal adenomas with sulindac in familial adenomatous polyposis.** *N Engl J Med* 1993, **328**:1313-1316.
- Giovannucci E, Egan KM, Hunter DJ, Stampfer MJ, Colditz GA, Willett WC, Speizer FE: **Aspirin and the risk of colorectal cancer in women.** *N Engl J Med* 1995, **333**:609-614.
- Schreinemachers DM, Everson RB: **Aspirin use and lung, colon, and breast cancer incidence in a prospective study.** *Epidemiology* 1994, **5**:138-146.
- Sharpe CR, Collet JP, McNutt M, Belzile E, Boivin J-F, Hanley JA: **Nested case-control study of the effects of non-steroidal anti-inflammatory drugs on breast cancer risk and stage.** *Br J Cancer* 2000, **83**:112-120.
- Baron JA, Sandler RS, Bresalier RS, Quan H, Riddell R, Lanis A, Bolognese JA, Oxenius B, Horgan K, Loftus S, Morton DG, APPROVe Trial Investigators: **A randomized trial of rofecoxib for the chemoprevention of colorectal adenomas.** *Gastroenterology* 2006, **131**:1674-1682.
- Funk CD: **Prostaglandins and leukotrienes: advances in eicosanoid biology.** *Science* 2001, **294**:1871-1875.
- Bennett A, McDonald AM, Stamford IF, Charlier EM, Simpson JS, Zebro T: **Prostaglandins and breast cancer.** *Lancet* 1977, **310**:624-626.
- Smith CJ, Sun D, Hoegler C, Roth BS, Zhang X, Zhao G, Xu XB, Kobari Y, Pritchard K Jr, Sessa WC, Hintze TH: **Reduced gene expression of vascular endothelial NO synthase and cyclooxygenase-1 in heart failure.** *Circ Res* 1996, **78**:58-64.
- Smith WL, DeWitt DL, Garavito RM: **Cyclooxygenases: structural, cellular, and molecular biology.** *Annu Rev Biochem* 2000, **69**:145-182.
- Kujubu DA, Fletcher BS, Varnum BC, Lim RW, Herschman HR: **TIS10, a phorbol ester tumor promoter-inducible mRNA from Swiss 3T3 cells, encodes a novel prostaglandin synthase/cyclooxygenase homologue.** *J Biol Chem* 1991, **266**:12866-12872.
- Xie WL, Chipman JG, Robertson DL, Erikson RL, Simmons DL: **Expression of a mitogen-responsive gene encoding prostaglandin synthase is regulated by mRNA splicing.** *Proc Natl Acad Sci USA* 1991, **88**:2692-2696.
- Soslow RA, Dannenberg AJ, Rush D, Woerner BM, Khan KN, Masferrer J, Koki AT: **COX-2 is expressed in human pulmonary, colonic, and mammary tumors.** *Cancer* 2000, **89**:2637-2645.
- Liu CH, Chang SH, Narko K, Trifan OC, Wu MT, Smith E, Haudenschild C, Lane TF, Hla T: **Overexpression of cyclooxygenase-2 is sufficient to induce tumorigenesis in transgenic mice.** *J Biol Chem* 2001, **276**:18563-18569.
- Tavassoli FA, Devilee P: *Pathology and Genetics of Tumours of the Breast and Female Genital Organs* Lyon: IARC Press; 2003.
- Sobin LH, Wittekind C: *TNM Classification of Malignant Tumours*. 6 edition. New York: Wiley-Liss; 2002.
- Elston EW, Ellis IO: **Method for grading breast cancer.** *J Clin Pathol* 1993, **46**:189-190.
- ArrayExpress. [http://www.ebi.ac.uk/arrayexpress].
- Dai M, Wang P, Boyd AD, Kostov G, Athey B, Jones EG, Bunney WE, Myers RM, Speed TP, Akil H, Watson SJ, Meng F: **Evolving gene/transcript definitions significantly alter the interpretation of GeneChip data.** *Nucleic Acids Res* 2005, **33**:e175.
- Akaike H: **Information theory and an extension of the maximum likelihood principle.** In *Second International Symposium of Inference Theory*. Edited by: Petrov B, Csaki F. Budapest: Akadémiai Kiadó; 1973:267-281.
- Ihaka R, Gentleman R: **R: a language for data analysis and graphics.** *Journal of Computational and Graphical Statistics* 1996, **5**:299-314.
- Lindsey J: *Models for Repeated Measurements*. 2 edition. Oxford: Oxford University Press; 1999.
- Dennis G, Sherman B, Hosack D, Yang J, Gao W, Lane HC, Lempicki R: **DAVID: Database for Annotation, Visualization, and Integrated Discovery.** *Genome Biol* 2003, **4**:3.
- van Iersel M, Kelder T, Pico A, Hanspers K, Coort S, Conklin B, Evelo C: **Presenting and exploring biological pathways with PathVisio.** *BMC Bioinformatics* 2008, **9**:399.
- Pico AR, Kelder T, van Iersel MP, Hanspers K, Conklin BR, Evelo C: **WikiPathways: pathway editing for the people.** *PLoS Biol* 2008, **6**:e184.
- Harper-Wynne C, Ross G, Sacks N, Salter J, Nasiri N, Iqbal J, A'Hern R, Dowsett M: **Effects of the aromatase inhibitor letrozole on normal breast epithelial cell proliferation and metabolic indices in postmenopausal women: a pilot study for breast cancer prevention.** *Cancer Epidemiol Biomarkers Prev* 2002, **11**:614-621.
- Kaiser CL, Chapman BJ, Guidi JL, Terry CE, Mangiardi DA, Cotanche DA: **Comparison of activated caspase detection methods in the gentamicin-treated chick cochlea.** *Hear Res* 2008, **240**:1-11.
- da Silva BB, Pires CG, dos Santos AR, Castro-Leão AHD, Alencar AP, Lopes-Costa PV: **Effects of raloxifene on Ki-67 and CD34 antigen expression in breast cancer.** *Gynecol Obstet Invest* 2009, **67**:103-108.
- Chalkley H: **Method for the quantitative morphologic analysis of tissues.** *J Natl Cancer Inst* 1943, **4**:47-53.
- Sato H, Kinoshita T, Takino T, Nakayama K, Seiki M: **Activation of a recombinant membrane type 1-matrix metalloproteinase (MT1-MMP) by furin and its interaction with tissue inhibitor of metalloproteinases (TIMP)-2.** *FEBS Lett* 1996, **393**:101-104.
- Talmadge JE: **Pathways mediating the expansion and immunosuppressive activity of myeloid-derived suppressor cells and their relevance to cancer therapy.** *Clin Cancer Res* 2007, **13**:5243-5248.
- Callegari AJ, Kelly TJ: **Shedding light on the DNA damage checkpoint.** *Cell Cycle* 2007, **6**:660-666.
- Dvory-Sobol H, Cohen-Noyman E, Kazanov D, Figer A, Birkenfeld S, Madar-Shapiro L, Benamouzig R, Arber N: **Celecoxib leads to G2/M arrest by induction of p21 and down-regulation of cyclin B1 expression in a p53-independent manner.** *Eur J Cancer* 2006, **42**:422-426.
- Liu W, Chen Y, Wang W, Keng P, Finkelstein J, Hu D, Liang L, Guo M, Fenton B, Okunieff P, Ding I: **Combination of radiation and celebrex (celecoxib) reduce mammary and lung tumor growth.** *Am J Clin Oncol* 2003, **26**:103-109.
- Dittmann KH, Mayer C, Ohneseit PA, Raju U, Andrantschke NH, Milas L, Rodemann HP: **Celecoxib induced tumor cell radiosensitization by inhibiting radiation induced nuclear EGFR transport and DNA-repair: a COX-2 independent mechanism.** *Int J Radiat Oncol Biol Phys* 2008, **70**:203-212.
- Fatima N, Yi M, Ajaz S, Stephens RM, Stauffer S, Greenwald P, Munroe DJ, Ali IU: **Altered gene expression profiles define pathways in colorectal cancer cell lines affected by celecoxib.** *Cancer Epidemiol Biomarkers Prev* 2008, **17**:3051-3061.

42. Auman JT, Church R, Lee SY, Watson MA, Fleshman JW, McLeod HL: **Celecoxib pre-treatment in human colorectal adenocarcinoma patients is associated with gene expression alterations suggestive of diminished cellular proliferation.** *Eur J Cancer* 2008, **44**:1754-1760.
43. Orlichenko L, Radisky D: **Matrix metalloproteinases stimulate epithelial-mesenchymal transition during tumor development.** *Clin Exp Metastasis* 2008, **25**:593-600.
44. Wang S, Liu Q, Zhang Y, Liu K, Yu P, Liu K, Luan J, Duan H, Lu Z, Wang F, Wu E, Yagasaki K, Zhang G: **Suppression of growth, migration and invasion of highly-metastatic human breast cancer cells by berbamine and its molecular mechanisms of action.** *Mol Cancer* 2009, **8**:81.
45. Wang L, Chen W, Xie X, He Y, Bai X: **Celecoxib inhibits tumor growth and angiogenesis in an orthotopic implantation tumor model of human colon cancer.** *Exp Oncol* 2008, **30**:42-51.
46. Lang S, Lauffer L, Clausen C, Lohr I, Schmitt B, Holzel D, Wollenberg B, Gires O, Kastenbauer E, Zeidler R: **Impaired monocyte function in cancer patients: restoration with a cyclooxygenase-2 inhibitor.** *FASEB J* 2003, **17**:286-288.
47. Lonnroth C, Andersson M, Arvidsson A, Nordgren S, Brevinge H, Lagerstedt K, Lundholm K: **Preoperative treatment with a non-steroidal anti-inflammatory drug (NSAID) increases tumor tissue infiltration of seemingly activated immune cells in colorectal cancer.** *Cancer Immun* 2008, **8**:5.
48. Harper-Wynne CL, Sacks NP, Shenton K, MacNeill FA, Sauven P, Laidlaw IJ, Rayer Z, Miall S, Howes A, Salter J, Hills MJ, Lowe FM, A'Hern R, Nasiri N, Doody D, Iqbal J, Dowsett M: **Comparison of the systemic and intratumoral effects of tamoxifen and the aromatase inhibitor vorozole in postmenopausal patients with primary breast cancer.** *J Clin Oncol* 2002, **20**:1026-1035.
49. Smith IE, Dowsett M, Ebbs SR, Dixon JM, Skene A, Blohmer JU, Ashley SE, Francis S, Boeddinghaus I, Walsh G: **Neoadjuvant treatment of postmenopausal breast cancer with anastrozole, tamoxifen, or both in combination: the Immediate Preoperative Anastrozole, Tamoxifen, or Combined with Tamoxifen (IMPACT) multicenter double-blind randomized trial.** *J Clin Oncol* 2005, **23**:5108-5116.
50. Brueggemeier R, Quinn A, Parrett M, Joarder F, Harris R, Robertson F: **Correlation of aromatase and cyclooxygenase gene expression in human breast cancer specimens.** *Cancer Lett* 1999, **140**:27-35.
51. Dowsett M, Ebbs SR, Dixon JM, Skene A, Griffith C, Boeddinghaus I, Salter J, Detre S, Hills M, Ashley S, Francis S, Walsh G, Smith IE: **Biomarker changes during neoadjuvant anastrozole, tamoxifen, or the combination: influence of hormonal status and HER-2 in breast cancer - a study from the IMPACT trialists.** *J Clin Oncol* 2005, **23**:2477-2492.
52. Dowsett M, Bundred NJ, Decensi A, Sainsbury RC, Lu Y, Hills MJ, Cohen FJ, Veronesi P, O'Brien ME, Scott T, Muchmore DB: **Effect of raloxifene on breast cancer cell Ki67 and apoptosis: a double-blind, placebo-controlled, randomized clinical trial in postmenopausal patients.** *Cancer Epidemiol Biomarkers Prev* 2001, **10**:961-966.
53. Martin LA, Davies G, Weigel M, Betambeau N, Hills M, Salter J, Walsh G, A'Hern R, Dowsett M: **Pre-surgical study of the biological effects of the selective cyclo-oxygenase-2 inhibitor celecoxib in patients with primary breast cancer.** *Breast Cancer Res Treat* 2010, **123**:829-836.
54. Subbaramaiah K, Norton L, Gerald W, Dannenberg AJ: **Cyclooxygenase-2 is overexpressed in HER-2/neu-positive breast cancer: evidence for involvement of AP-1 and PEA3.** *J Biol Chem* 2002, **277**:18649-18657.
55. Dang CT, Dannenberg AJ, Subbaramaiah K, Dickler MN, Moasser MM, Seidman AD, D'Andrea GM, Theodoulou M, Panageas KS, Norton L, Hudis CA: **Phase II study of celecoxib and trastuzumab in metastatic breast cancer patients who have progressed after prior trastuzumab-based treatments.** *Clin Cancer Res* 2004, **10**:4062-4067.
56. Dirix LY, Ignacio J, Nag S, Bapsy P, Gomez H, Raghunadharao D, Paridaens R, Jones S, Falcon S, Carpentier M, Abbattista A, Lobelle JP: **Treatment of advanced hormone-sensitive breast cancer in postmenopausal women with exemestane alone or in combination with celecoxib.** *J Clin Oncol* 2008, **26**:1253-1259.
57. Falandry C, Canney PA, Freyer G, Dirix LY: **Role of combination therapy with aromatase and cyclooxygenase-2 inhibitors in patients with metastatic breast cancer.** *Ann Oncol* 2009, **20**:615-620.
58. Algra AM, Rothwell PM: **Effects of regular aspirin on long-term cancer incidence and metastasis: a systematic comparison of evidence from observational studies versus randomised trials.** *Lancet Oncol* 2012, **13**:518-527.
59. Rothwell PM, Price JF, Fowkes FG, Zanchetti A, Roncaglioni MC, Tognoni G, Lee R, Belch JF, Wilson M, Mehta Z, Meade TW: **Short-term effects of daily aspirin on cancer incidence, mortality, and non-vascular death: analysis of the time course of risks and benefits in 51 randomised controlled trials.** *Lancet* 2012, **379**:1602-1612.
60. Snead D, Bell J, Dixon A, Nicholson R, Elston C, Blamey R, Ellis I: **Methodology of immunohistological detection of oestrogen receptor in human breast carcinoma in formalin-fixed, paraffin-embedded tissue: a comparison with frozen section methodology.** *Histopathology* 1993, **23**:233-238.

doi:10.1186/bcr3409

Cite this article as: Brandão et al.: A randomised controlled phase II trial of pre-operative celecoxib treatment reveals anti-tumour transcriptional response in primary breast cancer. *Breast Cancer Research* 2013 **15**:R29.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

